


Recombinant Protein G Conjugated to Sepharose Resin

Catalog No.	■	LT12015
Packing Details	■	1 mL, crosslinked 6% beaded agarose supplied as 50% slurry
Binding Capacity	■	20-25 mg human IgG per mL of settled resin
Support pH Stability	■	2-14 (short term); 3-13 (long term)
Particle Size	■	45 to 165 microns
Volumetric Flow	■	Approx. 1mL/minute ; Maximum Linear Velocity: 30cm per hour; Maximum Pressure: less than 4.5psi (0.3 bar)
Storage	■	4 °C - 8 °C in 20% ethanol
Shelf Life	■	3 years



Protein G Binding buffer: 20 mM sodium phosphate, 150 mM NaCl, pH 7.4
 Protein G Elution buffer: 100 mM Glycine, pH 3.0
 Protein G Neutralization buffer: 1M Tris buffer, pH 7.5-9

Protocol

1. Wash the prepacked column with 5~10 column volumes of distilled water to remove 20% ethanol.
2. Equilibrate the column with 5~10 column volumes of binding buffer.
3. 1:10 dilution of serum with binding buffer. Filtrate the diluted serum through a 0.45 µm filter and load the sample.
4. Wash with 10 column volumes of binding buffer.
5. Elute with 5 column volumes of elution buffer and neutralize collect fractions with neutralization buffer.
6. After each separation cycle, regenerate the resin by washing with approximately 3~5 column volumes of 0.1 M citrate buffer (pH 3.0).
7. Confirm the purity of the collected antibody by SDS-PAGE analysis.
8. Purification capacity: ≥30 mg of rabbit IgG per ml of recombinant protein G agarose gel.

Notes

1. Avoid air bubbles.
2. Regenerate the resin every 5 purification in order to maintain the product efficiency.
3. After every 10 separation cycle, wash the resin with 5 column volumes of 20% ethanol and 5~10 column volumes of 70% ethanol sequentially to remove hydrophobic substances.
4. Keep at 4~8 °C in 20% ethanol.

For research use only!